







Cystic Degeneration of the Peroneus Brevis Tendon: Ultrasound Diagnosis

Degeneração Quística do Tendão Fibular Curto: Diagnóstico Ecográfico

Bruno Werner Rezende Casimiro ¹, Denise de Aquino Gomes ¹, Gabriella de Macedo Calabrese ¹, Dalmo José Gonçalves Netto Borges ¹, Márcio Luís Duarte ^{2*}, Élcio Roberto Duarte ³

*Corresponding Author/Autor Correspondente

Márcio Luís Duarte [marcioluisduarte@gmail.com]

Universidade de Ribeirão Preto, Campus Guarujá. Av. D. Pedro I, 3.300, Enseada, Guarujá-SP, Brazil, ZIP CODE: 11440-003.

ORCID: <https://orcid.org/0000-0002-7874-9332>

<https://doi.org/10.48687/lj.203>

Keywords: Ganglion Cysts/diagnostic imaging; Tendon Injuries; Tendons/diagnostic imaging; Ultrasonography

Palavras-chave: Ecografia; Lesões dos Tendões; Quistos Ganglionares/diagnóstico por imagem; Tendões/diagnóstico por imagem

A 62-year-old female patient presented with complaints of bulging and pain in the lateral region of her foot for the past three months. She reported progressive bulging but noted no foot movement or shoe-wearing limitations. Upon physical examination, bulging was observed in the region of the base of the fifth metatarsal, extending to the hindfoot, without evidence of local hematomas or alterations in mobility. Palpation revealed that the bulging was mobile and hardened. Ultrasound examination revealed extensive septated cystic formations in the peroneus brevis tendon, suggesting cystic degeneration (Fig. 1). The patient underwent conservative treatment with rest and lifestyle changes in the subsequent weeks, resulting in an improvement in pain.

Ganglion or synovial cysts are benign tumours that can affect any area of the connective tissue of bones, muscles, and tendons. They are more common in women aged 15 to 40 years. Although ganglion cysts are recurrent, cysts originating from tendons, known as intratendinous ganglion cysts, are rare, especially when affecting the peroneal tendon.¹ Peroneal or

fibular tendons serve as lateral stabilizers of the ankle and can present various anatomical variations, making them prone to injuries. However, injuries to the peroneal tendons are often overlooked when patients complain of lateral ankle pain. Recognising the origin of this pain is essential to prevent future ankle dysfunctions and ensure a safe return to the patient's daily activities.²

For the evaluation of peroneal tendons, magnetic resonance imaging (MRI) is considered the gold standard examination, as it allows for better visualization of soft tissues in different planes, such as axial, sagittal, and coronal. Other imaging modalities, such as ultrasound, may be used for complementary purposes or in cases where MRI is contraindicated. In intratendinous ganglion cysts, MRI typically reveals a well-defined cystic lesion within the tendon, exhibiting high signal intensity on T2-weighted images and peripheral enhancement with contrast. MRI can help determine the extent of the lesion and assist in surgical planning.² Ultrasound, as demonstrated in this case, can also be used to diagnose intratendinous cysts,

1. Faculdade de Ciências Médicas de Santos, Santos, Brazil. **2.** Universidade de Ribeirão Preto – Campus Guarujá, Guarujá, Brazil. **3.** Irmandade da Santa Casa de Misericórdia de Santos, Santos, Brazil.

Recebido/Received: 05/04/2024 – **Aceite/Accepted:** 03/06/2024 – **Publicado online/Published online:** 12/06/2024 – **Publicado / Published:** 28/06/2024

© Author(s) (or their employer(s)) and Lusíadas Scientific Journal 2024. Re-use permitted under CC BY-NC 4.0. No commercial re-use.

© Autor (es) (ou seu (s) empregador (es)) e Lusíadas Scientific Journal 2024. Reutilização permitida de acordo com CC BY-NC 4.0. Nenhuma reutilização comercial.

offering advantages in terms of speed and cost compared to MRI.

Once the diagnosis is confirmed, treatment depends on two factors: the duration of signs and symptoms (acute or chronic) and whether ankle eversion is compromised. In simpler cases, conservative measures such as immobilization of the involved joint, rest, and lifestyle changes are often sufficient.^{1,2} Puncture

of the cyst or the use of injectable anti-inflammatory agents may be considered as an alternative treatment option. Surgical intervention is indicated for unstable and symptomatic ganglion cysts with severe and chronic characteristics.^{1,2} Surgical treatment involves not only excision of the cyst but also thorough examination and potential correction of underlying anatomical anomalies and fractures, as well as tendon debridement, tubularization, tenodesis, or autograft.¹




Figure 1. Ultrasound in longitudinal section (A), with Doppler (B) and in the axial section (C) demonstrating peroneus brevis tendon (black arrows) cystic degeneration (white arrows).

Responsabilidades Éticas

Conflitos de Interesse: Os autores declaram a inexistência de conflitos de interesse na realização do presente trabalho.

Fontes de Financiamento: Não existiram fontes externas de financiamento para a realização deste artigo.

Confidencialidade dos Dados: Os autores declaram ter seguido os protocolos da sua instituição acerca da publicação dos dados de doentes.

Consentimento: Consentimento do doente para publicação obtido.

Proveniência e Revisão por Pares: Não comissionado; revisão externa por pares.

Ethical Disclosures

Conflicts of Interest: The authors have no conflicts of interest to declare.

Financing Support: This work has not received any contribution, grant or scholarship.

Confidentiality of Data: The authors declare that they have followed the protocols of their work center on the publication of data from patients.

Patient Consent: Consent for publication was obtained.

Provenance and Peer Review: Not commissioned; externally peer reviewed.

Contributorship Statement

BWRC, DAG, GMC, MLD and ERD: Data curation, formal analysis, research, methodology, original draft writing, writing - proof-reading and editing.

All authors approved the final version to be published.

Declaração de Contribuição

BWRC, DAG, GMC, MLD e ERD: Curadoria de dados, análise formal, pesquisa, metodologia, redação do rascunho original, redação - revisão e edição.

Todos os autores aprovaram a versão final a ser publicada.

References

1. Frangež I, Kos TN, Cimerman M. Intratendinous ganglion cyst of peroneus brevis tendon and its reconstruction with semitendinosus graft. *Vojnosanit Pregl.* 2017; 74: 896–9.
2. Lima BF, Castro M, Costa AS, Brito N, Madureira AJ. Magnetic resonance imaging of common and uncommon peroneal tendons disorders. *Act Radiol Port.* 2022;34:9-16.
3. Robertson DE. Cystic degeneration of the peroneus brevis tendon. *J Bone Joint Surg Br.* 1959;41-B:362-4.